

RELATION BETWEEN PERITONEAL AGGRESSION AND FIBRINOLYSIS

Sebastian GRADINARU FOMETESCU¹, Costel SAVLOVSCHI¹,
 Marieta COSTACHE², Sorin Mircea OPRESCU¹

¹ University Emergency Hospital Bucharest, 4th Department of Surgery, Romania

² University of Bucharest, Department of Biochemistry and Molecular Biology, Romania

ABSTRACT. The triggering of the formation of adhesions is caused by the organization of the fibrin matrix, an organization that takes place during the coagulation processes when there is suppression of fibrinolysis. In this study two groups of patients are analyzed First group was used, both for evidencing the peritoneal fibrinolytic activity by determining the fibrinogen and the fibrin degradation products (FDP) in the peritoneal liquid and for extract the tPA and PAI-1 from the peritoneal tissue. The second group is the control group in whose cases none of the aggression factors (traumatic, chemical, bacterial) noticed in the patients of the first group was proven to exist. The aim of this study was to evaluate the role of PAI-1 and tPA genes at peritoneal tissue level. Peritoneal tissue was obtained during surgery and we have quantified the variation of expression of PAI-1 and tPA genes. The obtained results highlight an increase of expression in PAI-1 gene and decrease of expression in tPA gene, which indicated an decrease fibrinolytic potential.

Keywords: adhesions, fibrinolysis, fibrin, tPA, PAI-1

INTRODUCTION

The formation of adherences following abdominal and pelvic operations remains still frequent enough and is an important source of morbidity. The incidence of intraperitoneal adhesions ranges from 67% to 93% after general abdominal operations and up to 97% after gynecological operations in the pelvic region. The factors associated with the formation of postsurgical adhesions include: trauma, thermal injuries, infection, ischemia, and alien bodies.¹⁵⁷ Many other factors, including very tight sutures, when the tension in the sutured peritoneum brings about ischemia, abrasion, exposing to aggression factors such as talcum powder in the gloves, reactive sutures materials, contents of bowels, superheating due to the lamps or to the irrigation fluid, can all contribute to the formation of postsurgical adhesions.

The endometriosis, one of the most frequent causes of pelvic pain or infertility in women at the optimum age of reproduction, is associated with the formation of fibrous adherences following operations with the preservation of organs and operations of excision of endometria. Fibrous adhesions often form as a response to the chronic irritation of the peritoneal surface caused by the endometric implant and its secretion products.

The formation of adherences occurs, typically, in the place, where two peritoneal surfaces come in contact. The triggering of the formation of adhesions is caused by the organization of the fibrin matrix, an organization that takes place during the coagulation processes when there is suppression of fibrinolysis. The tissular surgical lesions diminish or eliminate the blood flow, giving rise to ischemia the effect of which being the local persistence of the fibrous matrix, gradually replaced vascular tissue of granulation, which contains macrophages, fibroblasts, and giant

cells. The adhesions can subsequently mature in the shape of fibrous bands, which often contain small calcified nodules and are frequently covered with mesothelium, blood vessels and fibers of conjunctive tissue, inclusively elastin.

MATERIALS AND METHODS

Structure of patient groups. In this study two groups of patients are analyzed.

The **first group** is composed of twelve patients operated on in the clinic for surgical acute abdomen, in whose case the presence of a factor of peritoneal aggression (FPA) susceptible to bring about the formation of postsurgical peritoneal adhesions was particularly noticeable during the operation. The factors of peritoneal aggression have been divided in three categories: traumatic, chemical, and bacterial. This group was used, both for evidencing the peritoneal **fibrinolytic activity** by determining the fibrinogen and the fibrin degradation products (FDP) in the peritoneal liquid and for extract the tPA and PAI-1 from the peritoneal tissue.

The **second group** is the control group made up of six patients operated on in the clinic for defects of the abdominal wall (three patients) and for acute catarrhal appendicitis with peritoneal serous reaction (three patients), respectively, taking into consideration the fact that – in these cases – none of the aggression factors noticed in the patients of the first group was proven to exist. The patients have consented to be included in this study, which was approved by the ethical committee of the hospital.

Processing of samples of peritoneal liquid with the view to making salient the fibrinolytic activity

For the patients in the first group, the peritoneal liquid was collected by aspiration from the drainage

tube located as near to the seat of the maximum peritoneal aggression as possible, beginning from the first day after the operation and repeating the collection daily up to the seventh day. For determining the concentration of fibrinogen, the peritoneal liquid was processed in the emergency laboratory of the Emergency University Hospital of Bucharest, where it was sent in a vacuum container for the coagulogram to be made. For the quantitative determination of the fibrin degradation products, the peritoneal liquid was collected in a syringe containing 0.1 ml of heparin and was processed on a Compact Immuno-analyser Pathfast Sintergy (Mitsubishi Kagaku Iatron.Inc), utilizing Pathfast D-dimer (Mitsubishi Kagaku Iatron.Inc) kits.

For the patients in the witness group, the peritoneal liquid was first collected during the operation and afterwards daily – for two or three days – from a thin tube located in the bottom of the Douglas sac, until the drainage diminishes and the drainage tube is taken out.

Processing of the peritoneal tissue with the view to determining tPA and PAI-1

In order to extract the tPA and PAI-1 from the peritoneal tissue, fragments of peritoneal tissues freshly collected during the operation from the patients operated on for surgical acute abdomen were used, in which the presence of one of the peritoneal aggression factors indicated in the table below had been established. For comparison purposes, the witness group was made up of patients with uncomplicated hernias of the abdominal wall, considering that – at the level of the sacular peritoneum – none of the studied aggression factors act, and the collection of peritoneal tissue does not involve any supplementary surgical gestures since the resection of the hernia sac or the eventration represent a usual stage within the framework of the surgical treatment of the abdominal parietal defects.

Diagnosis	Traumatic FPA	Chemical FPA	Bacterial FPA
Hemoperitoneum by spleen rupture	blood	0	0
Hemoperitoneum by entero-mesenteric wound	blood	0	0
Perforated duodenal ulcer	0	bile+gastric juice	0
Perforated gastric ulcer	0	gastric juice	0
Perforated gangrenous calculous cholecystitis	0	infected bile	<i>E. coli</i>
Biliary septic peritonitis	0	uninfected bile	0
Perforated gangrenous acalculous cholecystitis	0	0	<i>E. coli</i>
Biliary chemical peritonitis	0	0	<i>E. coli</i>
Perforated gangrenous appendicitis	0	urine+blood	0
Generalized peritonitis	0	infected urine	<i>E. coli</i>
Diastatic caecum perforation due to colon cancer.	0	endometriotic liquid	0
Generalized peritonitis	0	0	<i>E. coli</i>
Uroperitoneum by traumatic rupture of bladder	0	0	0
Uroperitoneum by PNL perforation of renal pelvis	0	0	<i>E. coli</i>
Ruptured ovarian endometriotic cyst	0	0	0
Endometriotic peritonitis	0	0	<i>E. coli</i>
Ruptured tuboovarian abscess. Pelvipertonitis	0	0	<i>E. coli</i>

Biopsy procedure

The biopsy was conducted on the parietal peritoneum (punch for biopsy with a 6-mm diameter, available, Stiefel Lab., Woobum Green, UK) as quickly as possible after the opening of the peritoneal cavity. The tissue disk was dissected from the adjacent subperitoneal tissue and minced in fragments with dimensions smaller than 1 mm, which were introduced in recipients covered with RNA stabilizer and kept in a refrigerator at a temperature of 4°C until they were processed.

Extraction of RNA from the tissue using the RNeasy® LIPID TISSUE KIT

Principle of method

The RNeasy Lipid Tissue Kit combines the method of lysis of the cells on the basis of phenol-guanidine with purification of the total RNA by means of minicolumns of silicon dioxide. The lysis reactive QIAzol is a monophasic solution of phenol and

guanidine thiocyanate, ideal for the lysis of the various tissues (mainly in the case of tissues containing much fat) and for the inhibition of RNase. In a first stage, the following operations are conducted: homogenization of the tissue in the lysis reactive QIAzol, addition of chloroform and separation – by centrifuging – of the homogenate in the aqueous phase and the organic phase. In the following stage, the superior phase, which is aqueous (and contains the total RNA) is collected and then mixed with ethanol and the mixture is passed through an RNeasy minicolumn. The total RNA is then tied to the little column, and the phenol – as well as other contaminants – are eliminated. After successive washing by means of various washing swabs, the total RNA is eluted in water without RNase.

Statistical evaluation of results

The interpretation of results was effected by utilizing the statistical analysis conducted by means of the Excel and Epi Info computer programs. The

homogeneity of data distribution was determined by the Bartlett test. A value of p lower than 0.05 of the Bartlett test is normally considered to be significant and it shows a non-homogenous distribution of the data to be analyzed, while a value of p equal to or higher than 0.05 indicates a similar dispersion of the subsamples. The significance of the differences between the patient groups with peritoneal risk factors and without peritoneal risk factors was determined by utilizing the parametric tests Student for two groups and ANOVA for more than two groups, as well as the non-parametric tests Mann-Whitney for two groups and Kruskal-Wallis for more than two groups. The difference is statistically significant if the probability p of these tests is lower than 0.05.

The Student and ANOVA tests are applicable only in the case of a homogenous data distribution (the Bartlett test has a $p \geq 0.05$). In the case of a non-homogenous dispersion of subsamples (the Bartlett test has a $p < 0.05$), the Bartlett test is not appropriate and a non-parametric statistics (the Mann-Whitney and Kruskal-Wallis tests) is to be used.




Fig. 1 Temperature gradient for optimizing the amplification conditions of a fragment of PAI-1 gene.
1: 51°C; 2: 51,8°C; 3: 53,1°C; 4: 54,9°C; 5: 57,4°C; 6: 59,3°C; 7: 60°C; 8: 50bp molecular weight marker

Figure 3 shows the amplification curves which represent the fluorescent signals obtained during the reaction. In fact the graph represents the intensity of the fluorochrome interpolated at the level of the PCR products (thus measuring, in the case of the same gene, the quantity of PCR product formed in each cycle) vs the amplification cycle.

To calculate the efficiency of a PCR reaction, a logarithmic graph is obtained in which the values of C_t are represented vs the dilutions of the sample. In Figs. 4 and 5, the efficiencies of the PCR reaction are represented for the case of amplification of fragments of PAI-1 and actin genes.

The efficiency of the PCR reaction is considered to be good if it is 90-110%, which corresponds to a slope

RESULTS

Analysis of peritoneal-tissue samples. Relative quantification of the expression of PAI-1 gene

In order to study the expression of PAI-1 gene, the Real-Time PCR procedure was used, going through the following stages: reverse transcription of total RNA into cDNA, optimization of amplification conditions (PAI-1 gene and reference gene), achievement of the standard curve and calculation of the efficiency of PCR reaction. Finally, all the samples were run and normalized as a function of the actin reference gene, and in the end the relative expression of PAI-1 was compared between various patients.

The PCR conditions were optimized for the primer by varying the annealing temperature between 50-60°C on a gradient thermocycler IQCycler (BioRad). The PCR products were directly analyzed by electrophoresis in 2% agarose gels in 1X TAE buffer, stained with ethidium bromide, and visualized under UV light (Figs. 1 and 2). The optimum annealing temperature for all genes was 54°C.




Fig. 2 Temperature gradient for optimizing the amplification conditions of a fragment of actin gene.
1: 51°C; 2: 51,8°C; 3: 53,1°C; 4: 54,9°C; 5: 57,4°C; 6: 59,3°C; 7: 60°C; 8: molecular-mass marker 50bp

value between -3.58 and -3.10. As it can be noticed in Figs. 4 and 5 as well, the efficiencies obtained in the case of amplification of fragments of PAI-1 genes and actin were 106.6% (slope -3.173; $R^2=1$) and 105% (slope -3.208; $R^2=0.998$).

After the calculation of efficiencies, samples from various patients were run in order to quantify the relative expression of the PAI-1 gene as a function of the reference gene – actin. In Fig. 6, the amplification curves for all samples are presented.

The obtained melting curves have shown the existence of a single peak (89°C – PAI-1; 84°C – actin), which means that there were no non-specific amplifications, dimers of primers or contaminations with genomic DNA.




Fig. 3 Standard curve was obtained using 5 serial dilutions of a cDNA sample: $10\times$ dilution (10ng/reaction), $10^2 \times$ dilution (1ng/reaction), $10^3 \times$ dilution (0.1ng/reaction), $10^4 \times$ dilution (0.01ng/reaction), $10^5 \times$ dilution (0.00ng/reaction)




Fig. 4 Efficiency of the amplification reaction of a fragment of PAI-1 gene




Fig. 5 Efficiency of the amplification reaction of a fragment of actin gene




Fig. 6 – Amplification curve in the case of quantification of the expression of PAI-1 gene.

The amplification equations were determined for both the target gene PAI-1 and the reference gene (actin); in the case of each sample we obtained the following equations:

$$N_{CtPAI-1} = N_{0PAI-1} (1+E)^{CtPAI-1},$$

$$N_{Ctactin} = N_{0actin} (1+E)^{Ctactin},$$

where

N_{Ct} = number of molecules after C_t amplification cycles,

N_0 = initial number of molecules,

E = efficiency of PCR reaction,

C_t = threshold cycle

For normalization purposes, the ratio between N_{0PAI-1} and N_0 of the reference gene (actin) was calculated with the formula:

$$N_{0PAI-1}/N_{0ref} = k 2^{C_{tref}-C_{tPAI-1}}$$

where k is a correction factor.

The above equation was used for all of the patients, including the normal patient (deemed to be the control patient). In accordance with the equation

$Control/patient = k 2^{C_{trefcontrol}-C_{tPAI-1control}}/k 2^{C_{trefpatient}-C_{tPAI-1patient}} = 2^{\Delta\Delta C_t}$, all of the samples were referred to the control patient, and the diagram in Fig. 7 was obtained.

In compliance with the statistical test t-test, two tails, non-paired, all of the values obtained for p were significant ($p < 0.05$). It is apparent from the diagram in Fig. 7 that the relative expression of PAI-1 gene is significantly increased (3-4.5 times) in comparison with the case of the sample considered control sample.

Relative quantification of the expression of tPA gene.

In order to study the relative expression of tPA gene, the Real-Time PCR technique was used as well. The actin gene was used as reference gene, and the stages run through were the same as in the case of the quantification of expression of PAI-1 gene: optimization of the hybridization temperature of primers, calculation of the efficiency of PCR reaction, and comparison of the expression of tPA of the patients.

The PCR conditions were optimized for the primer by varying the annealing temperature between 50-60°C on a gradient thermocycler IQCycler (BioRad). The PCR products were directly analyzed by electrophoresis in 2% agarose gels in 1X TAE buffer, stained with ethidium bromide, and visualized under UV light (Fig. 8).

The optimum temperature of the hybridization of primers was 54°C. In Fig. 9 we have shown the obtained standard curve for tPA gene. The efficiency of the amplification reaction of a fragment of tPA gene was 99%, with a slope of -3.346 and $R^2 = 0.998$.

After the calculation of efficiency, samples from various patients were run in order to quantify the relative expression of tPA gene as a function of the actin reference gene. The samples from each patient were run in triplicate (technical replications). After the samples were run, the program of the PCR instrument established the threshold level and calculated the values of C_t for each sample. In Fig. 10 the amplification curve of all of the samples is presented.




Fig. 7 – Relative expression PAI-1 gene




Fig. 8 Temperature gradient for the optimization of the amplification conditions of a fragment of tPA gene
 1: 51°C; 2: 51,8°C; 3: 53,1°C; 4: 54,9°C; 5: 57,4°C; 6: 59,3°C; 7: 60°C; 8: 50 bp molecular weight marker




Fig. 9 Efficiency of the amplification reaction of a fragment of tPA gene

The data obtained was processed in compliance with the equations given above, and the average values obtained – along with the standard deviation – were represented graphically in Fig. 11.

In compliance with the statistic test t-test, two tails, unpaired, all of the values obtained for p were significant ($p < 0.05$). In the case of tPA gene, a level of the expression was obtained, which was 3 to 9 times lower than in the case of the control patients (fig. 11).

Analysis of the samples of peritoneal liquid. Analysis of the fibrinogen/PDF balance in the samples of peritoneal liquid.

The samples were collected in a period of seven days after the operation from the twelve patients in the study lot, in which – during the operation – the presence of one of the aggression factors described in the table was ascertained. The samples from the patients in the witness lot were collected only in the first three days after the operation, as long as a drainage tube located in the Douglas sac bottom was kept.

The sample collection and the quantitative determinations were conducted in accordance with the protocol provided in the chapter "Materials and methods". The average age of the patients with factors of peritoneal aggression was 43 years (19 to 71 years),

and the average age of the patients in the witness lot was 29 years (24 to 32 years).

In view of the fact that the adherence formation of is a dynamic process, we found it necessary to collect the seriate samples from the same subjects, at different intervals from the surgical intervention, with dosage of inflammation and fibrinolysis markers in each stage of the study, this kind of research being able to offer much more important information regarding the evolution of physiological mechanisms.

The same variations of the studied parameters were found for all of the patients in the study lot, irrespective of the aggression factor involved. It is to be noted that the fibrinogen values exhibit a maximum in the fifth day after the operation, and afterwards they begin to decrease until the seventh day, however without reaching the minimum values noticed in the first day, but – in accordance with the conclusions of the previous study – the absolute values are directly proportional to the risk of adhesiogenesis.

The presence of fibrine d-dimers in the peritoneal liquid was expected since the fibrinogenesis entails sequentially the fibrinolysis processes, but this research was not intended to determine the extent to which FDP can be true indicators of the fibrinolytic activity and can correlate with the risk of adhesiogenesis.




Fig. 10 Amplification curve in the case of quantification of the expression of tPA gene






Fig. 11 Relative expression of tPA gene





In this respect, we notice that the average value of FDP correlates with the mean of the fibrinogen values for each individual case, but the quantitative variation of d-dimers is inversely proportional to the quantitative variation of fibrinogen until the fifth day after the operation and afterwards exhibits the same decreasing trend. These aspects can be found in the diagrams of synchronous variations.

One of the drawbacks of this study is the impossibility of comparing the results of the determinations made for the patients subjected to aggression factors with the results of the determinations made for the patients in the witness lot, for the whole period of seven days after the operation. For ethical and medical reasons, this statistical comparison was made only for the first 2 to 3 days after the operation. If for the patients in the study lot the collections of peritoneal liquid were possible for seven consecutive days, until the last drainage tube was removed from the peritoneal cavity, for the patients in the witness lot the collection of the intraperitoneal exudates was effected as long as the peritoneal

drainage was necessary for the patients with appendicitis and serous peritoneal reaction. The peritoneal liquid was collected from one of the patients in the control lot for three consecutive days and a descending trend of the values of FIB and FDP was noticed, which could suggest that the prolonged inflammation of the peritoneal serosa, under the influence of aggression factors, represents the main trigger of adhesiogenesis, although the fibrinolytic peritoneal activity was discovered in all of the patients in the study lot.

Unlike the fibrinogen case, the FDP values have a higher predictive character as regards the formation of peritoneal adherences, depending on the determining aggression factors. It was noticed that in the cases of anexial pathology (ruptured endometriotic cyst and pelviperitonitis by ruptured tuboovarian abscess), the FDP values are $1.7 \pm 1.104491 \text{ ng/ml}$ and $1.924286 \pm 0.984155 \text{ ng/ml}$ ($p < 0,05$) and they decrease very slowly, in the situation, where the quantity of fibrinogen in the peritoneal exudates and, implicitly, the resulting quantity of fibrin are the biggest

(3.859429±1.70499g/L and 3.384143±1.642043g/L, p<0,05) as compared with the other aggression factors, between their variations existing a negative correlation of -0.72239 and -0.69722 ($|r| < 1$), respectively.

An argument confirming this hypotesis is represented by the similar FDP values

(2.185714±0.571573ng/ml, p<0,05), but in the conditions of a much lower concentration of fibrinogen (2.614714±1.464473g/L, p<0,05) in the patients with post-traumatic hemoperitoneum.

Statistical Analysis	Fibrinogen (FIB) – g/L						
	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
(A) Patients with FPA							
-Average	0.8232	1.9999	2.3411	3.7773	5.3780	4.1944	3.1527
-Std. Dev.	0.4473	0.4056	0.5297	0.6692	0.6584	0.5237	0.3334
-Variation	0.2000	0.1645	0.2805	0.4478	0.4336	0.2742	0.1111
(B) Patients without FPA							
-Average	0.2080	0.5497					
-Std. Dev.	0.0316	0.0862					
-Variation	0.0010	0.0074					
(A)vs (B)							
Bartlett's Test	0.0073	0.0604					
P value							
ANOVA							
P-value		0.0000					
Kruskal-Wallis Test							
P value	0.0094						

Statistical Analysis	Fibrin degradation products (FDP) – ng/ml						
	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
(A) Patients with FPA							
- Average	4.0333	3.7067	3.3542	2.7567	2.2308	1.8117	1.2175
- Std. Dev.	1.3880	0.8861	0.9378	0.8168	0.9003	0.8676	0.7182
- Variation	1.9267	0.7852	0.8795	0.6672	0.8105	0.7528	0.5158
(B) Patients without FPA							
- Average	0.9000	0.8700					
- Std. Dev.	0.3517	0.2506					
- Variation	0.1237	0.0628					
(A)vs (B)							
Bartlett's Test	0.0852	0.1056					
P value							
ANOVA							
P-value	0.0023	0.0001					
Kruskal-Wallis Test							
P value							

Ord. No.	DIAGNOSIS	Correlation with control group $ r < 1$	FIB		Correlation FIB – PDF $ r < 1$
			g/l	t-test (p<0,05)	
1	Hemoperitoneum by spleen rupture	-0.16123	2.461429± 1.397516	0.007378	-0.59418
2	Hemoperitoneum by entero-mesenteric wound	-0.07954	2.614714± 1.464473	0.006611	-0.45705
3	Perforated duodenal ulcer	0.155893	2.764429± 1.429679	0.004435	-0.82322
4	Perforated gastric ulcer	0.292336	2.605±1.37 9216	0.014515	-0.72755
5	Perforated gangrenous calculous cholecystitis.	0.081281	3.158±1.52	0.002947	-0.61594

	Biliary septic peritonitis		0759		
6	Perforated gangrenous acalculous cholecystitis. Biliary chemical peritonitis	0.281678	2.933714± 1.481369	0.003901	-0.62173
7	Perforated gangrenous appendicitis Generalized peritonitis.	0.183364	2.865857± 1.587544	0.006133	-0.71399
8	Diastatic caecum perforation due to colon cancer. Generalized peritonitis.	-0.04347	2.994857± 1.557297	0.004443	-0.58184
9	Uroperitoneum by traumatic rupture of bladder.	0.410965	3.592429± 1.575757	0.001668	-0.73784
10	Uroperitoneum by PNL perforation of renal pelvis.	0.33738	3.908571± 1.570247	0.001	-0.65467
11	Ruptured ovarian endometriotic cyst. Endometriotic peritonitis.	-0.3222	3.859429± 1.70499	0.001644	-0.72239
12	Ruptured tuboovarian abscess. Pelvipertonitis.	-0.22008	3.384143± 1.642043	0.002794	-0.69722
M	Control group patients.		0.31781±0, 135661		0.482501

Ord. No.	DIAGNOSIS	Correlation with control group $ r < 1$	FDP ng/ml	t-test ($p < 0,05$)	Correlation FIB – PDF $ r < 1$
1	Hemoperitoneum by spleen ruptura	-0.16123	1.96±1.563 689	$p > 0,05$	-0.59418
2	Hemoperitoneum by entero-mesenteric wound	-0.07954	2.185714± 0.571573	0.00010184	-0.45705
3	Perforated duodenal ulcer	0.155893	2.337143± 1.423818	0.012572882	-0.82322
4	Perforated gastric ulcer	0.292336	2.687143± 1.42946	0.005745994	-0.72755
5	Perforated gangrenous calculous cholecystitis. Biliary septic peritonitis	0.081281	2.714286± 1.37902	0.004634731	-0.61594
6	Perforated gangrenous acalculous cholecystitis. Biliary chemical peritonitis	0.281678	3.32±1.003 594	0.000179397	-0.62173
7	Perforated gangrenous appendicitis Generalized peritonitis.	0.183364	3.081429± 0.831814	$p > 0,05$	-0.71399
8	Diastatic caecum perforation due to colon cancer. Generalized peritonitis.	-0.04347	2.558571± 0.822991	0.00032359	-0.58184

9	Uroperitoneum by traumatic rupture of bladder.	0.410965	4.37±1.000 95	p>0,05	-0.73784
10	Uroperitoneum by PNL perforation of renal pelvis.	0.33738	3.922857± 1.966814	0.003076069	-0.65467
11	Ruptured ovarian endometriotic cyst. Endometriotic peritonitis.	-0.3222	1.7±1.1044 91	0.026521573	-0.72239
12	Ruptured tuboovarian abscess. Pelvipertonitis.	-0.22008	1.924286± 0.984155	0.00677818	-0.69722
M	Control group patients.		0,658571± 0,16807		0.482501

DISCUSSIONS AND CONCLUSIONS

The physiological disappearance of the provisional matrix of the clot is as important as its formation. The main proteolytic enzymes, the activators of plasminogen and the plasmin avoid the action of such inhibitors of proteases as the inhibitor of the activator of plasminogen and the α 2-antiplasmin, linking themselves with the fibrine clot and the cellular surfaces. Although the activator of the plasminogen and the plasmin can degrade a wide range of proteins of the extracellular matrix, a specific inhibitor of the activator of plasminogen (PAI-1) links itself with the extracellular matrix and limits the matriceal degradation to a very small space neighbouring the cellular surfaces. The inadequate removal of fibrin can hinder the normal healing process of the wounds and may lead to the formation of fibrous adherences.

PAI-1 is a glycoprotein of 50kD produced at the level of the endothelial cells, which belongs to the superfamily of superprotease inhibitors (serpin-1) and represents the major inhibitor of the tPA activator (tissue plasminogen activator) and of the activator uPA (urokinase plasminogen activator), at the level of the plasma and of the peritoneum (Kruithof, 1988; van Hinsbergh *et al.*, 1990), determining the formation of the inactive enzymatic complexes PA-PAI. The high levels of expression of PAI-1 and tPA are associated with the coronarian risk at the level of arteries. The increased activity of tPA is an indicator of the increased fibrinolytic potential. The action of PAI-1 is controlled by various cytokines and growth factors, such as TGF- β . Thus, TGF- β stimulates the secretion of PAI-1 in various cellular lines (Rondeau *et al.*, 1995) and *in vivo* (Sawdey&Loskuttoff, 1991).

The fibrinolytic system plays an important role in the formation/reformation of the peritoneal adherence. In the first stages of the healing process, between two peritoneal surfaces wounded, from the fibrinous matrix fibrin bands develop subsequently, which can be removed in a natural way by fibrinolysis, the result being the formation of fibrin degradation products (FDP). In the pathological processes of the adherential

formation, the fibrinolytic mechanisms are alterate, permitting the formation and persistence of adherences.

In the light of the theoretical arguments, the quantitative determination of fibrinogen in the peritoneal liquid represents an indicator of the inflammation induced by the presence of aggression factors in the peritoneal cavity, while the quantitative determination of the products of degradation of fibrin (D-dimers) could be an indicator of the fibrinolytic activity.

In the case of formation or reformation of adherences, the most important period is the one composed of the first seven days after the operation, when all of the processes of sedimentation of fibrin and, subsequently, of fibrinolysis take place. The collection of the only sample of peritoneal liquid from the subjects with the intraperitoneal adherences already formed offers little information regarding the events of adherential formation that took place before the date of collection of the sample.

The continuation of research is necessary for knowing the mechanisms involved in adesiogenesis from a molecular and cellular point of view and for determining with more precision the role of fibrin, as well as the regulation of fibrinolysis and coagulation, all of these from the point of view of the involvement in the generation of adherences. Also, by understanding the biochemical and morphological phenomena of the normal peritoneal healing and of cicatrization, it will be possible to obtain a modality of separating the peritoneal surfaces and increasing fibrinolysis. The difference between the adherences that appeared at different intra-abdominal or intrapelvic levels must be studied further. A better understanding of the adherences that predispose to complications and the knowledge of the places where these appear more frequently is necessary. Further research is required for a better knowledge of the newly discovered agents, with special emphasis on the degree of adherence prevention. It is also necessary to identify the best diagnosis procedures to permit the visualization of the adherences that appeared after the operation.

REFERENCES

- Cheong YC, Laird SM, Li TC, et al. Peritoneal healing and adhesion formation/reformation. *Hum Reprod Update* 2001; 7:556-66.
- Ellis H. The magnitude of adhesion-related problems. *Ann Chir Gynaecol* 1998;87:9-11.
- Holmdahl L, Eriksson E, Al-Jabreen M, Risberg B. Fibrinolysis in human peritoneum during operation. *Surgery*. 1996;119(6):701-705.
- Imudia AN, Kumar S, Saed GM, Diamond MP. Pathogenesis of Intra-abdominal and pelvic adhesion development. *Semin Reprod Med*. 2008 Jul;26(4):289-97.
- Ivarsson M, Holmdahl L, Falk P, et al. Characterisation and fibrinolytic properties of mesothelial cells isolated from peritoneal lavage. *Scand J Clin Lab Invest* 1998;58: 195-203.
- Kruithof EKO (1988) Plasminogen activator inhibitors: A review. *Enzyme* 40:113-121
- Menzies D, Ellis H. Intestinal obstruction from adhesions--how big is the problem? *Ann R Coll Surg Engl* 1990; 72:60-63
- Moore KL, Bang NU, Brodie TA, et al. Peritoneal fibrinolysis: evidence for the efficiency of the tissue-type plasminogen activator. *J Lab Clin Med* 1983;101:921-9.
- Mutsaers SE, Prêle CM, Lansley SM, Herrick SE. The origin of regenerating mesothelium: a historical perspective. *Int J Artif Organs*. 2007 Jun;30(6):484-94.
- Pattinson H, Koninckx P, Brosens I, et al. Clotting and fibrinolytic activities in peritoneal fluid. *Br J Obstet Gynecol* 1981; 88: 160-6.
- Rodgers, K.E., and G.S. diZerega. 1993. Function of peritoneal exudate cells after abdominal surgery. *J. Invest. Surg.* 6:9-23.
- Rondeau E, Sraer JD, Schleuning WD (1995) The renal plasminogen activating system, in *Molecular Nephrology: Kidney Function in Health and Disease*, New York, Marcel Decker Inc., pp 699-715
- Sawdey MS, Loskuttoff DJ (1991) Regulation of murine type 1 plasminogen activator inhibitor gene expression in vivo. *J Clin Invest* 88:1346-1353
- Scott-Coombes DM, Whawell SA, Thompson JN. The operative peritoneal fibrinolytic response to abdominal operation. *Eur J Surg* 1995; 161: 395-399
- Sulaiman H, Dawson L, Laurent GJ, et al. Role of plasminogen activators in peritoneal adhesion formation. *Biochem Soc Trans* 2002;30:126-31.
- Thompson J, Paterson-Brown S, Harbourne T, et al. Reduced human peritoneal plasminogen activating activity: a possible mechanism of adhesion formation. *Br J Surg* 1989; 76: 382-84.
- van Hinsbergh VWM, Kooistra T, Scheffer A, van Bockel JH, van Muijen GNP (1990) Characterization and fibrinolytic properties of human omental tissue mesothelial cells: comparison with endothelial cells. *Blood* 75:1490-1497
- Vipond MN, Whawell SA, Thompson JN, et al. Peritoneal fibrinolytic activity and intra-abdominal adhesions. *Lancet* 1990; 335:1120-2.
- Vipond M, Whawell S, Thompson J, et al. Effect of experimental peritonitis and ischaemia on peritoneal fibrinolytic activity. *Eur J Surg* 1994; 160:471-77.
- Vrijland WW, Jeekel J, van Geldorp HJ, et al. Abdominal adhesions: intestinal obstruction, pain, and infertility. *Surg Endosc* 2003;17:1017-22
- Weibel MA, Manjo G. Peritoneal adhesions and their relationship to abdominal surgery. *Am J Surg* 1973;126: 345-53.